

IMMUNIZATION SCHEDULE

RECOMMENDED DURING PREGNANCY

Diseases prevented	Vaccine	During pregnancy
Tetanus, diphtheria & pertussis (Tdap)	Tdap vaccine	1 dose, preferably at 27–36 weeks' gestation*
Influenza (Flu)	Inactivated influenza vaccine or recombinant influenza vaccine**	Annual dose at any point during pregnancy
COVID-19 (Coronavirus disease 2019)	COVID-19 vaccine	COVID-19 vaccine, when recommended
Respiratory syncytial virus (RSV)	RSV vaccine	1 dose during pregnancy at 32–36 weeks' gestation for babies to be born during RSV season***

*If Tdap is administered before pregnancy, it should be administered again during pregnancy.

**Live attenuated influenza vaccine is not recommended during pregnancy.

***RSV season lasts from October through March in most of the continental United States; maternal RSV vaccine is given September–January. Women who received one dose of RSV vaccine during pregnancy should not receive additional doses during later pregnancies. Instead, that infant should receive nirsevimab.

These vaccines are recommended to protect you and to keep your baby safe, too!

Vaccine Information Statements: <https://www.cdc.gov/vaccines/hcp/vis/index.html>

Recommended Immunization Schedule for Adults
Aged 19 Years or Older — United States, 2024

